

We claim:

1. A method for simultaneous sequence-specific identification of mRNAs in a mRNA population comprising the steps of:

(a) preparing double-stranded cDNAs from a mRNA population using a mixture of 12 anchor primers, the anchor primers each including: (i) a tract of from 7 to 40 T residues; (ii) a site for cleavage by a restriction endonuclease that recognizes more than six bases, the site for cleavage being located to the 5'-side of the tract of T residues; (iii) a stuffer segment of from 4 to 40 nucleotides, the stuffer segment being located to the 5'-side of the site for cleavage by the restriction endonuclease; and (iv) phasing residues -V-N located at the 3' end of each of the anchor primers, wherein V is a deoxyribonucleotide selected from the group consisting of A, C, and G; and N is a deoxyribonucleotide selected from the group consisting of A, C, G, and T, the mixture including anchor primers containing all possibilities for V and N;

(b) producing cloned inserts from a suitable host cell that has been transformed by a vector, the vector having the cDNA sample that has been cleaved with a first restriction endonuclease and a second restriction endonuclease inserted therein, the cleaved cDNA sample being inserted in the vector in an orientation that is antisense with respect to a bacteriophage-specific promoter within the vector, the first restriction endonuclease recognizing a four-nucleotide sequence and the second restriction endonuclease cleaving at a single site within each member of the mixture of anchor primers;

(c) generating linearized fragments of the cloned inserts by digestion with at least one restriction endonuclease that is different from the first and second restriction endonucleases;

(d) generating a cRNA preparation of antisense cRNA transcripts by incubation of the linearized fragments with a bacteriophage-specific RNA polymerase capable of initiating transcription from the bacteriophage-specific promoter;

(e) dividing the cRNA preparation into sixteen subpools and transcribing first-strand cDNA from each subpool, using a thermostable reverse transcriptase and one of sixteen primers whose 3'-terminus is -N-N, wherein N is one of the four deoxyribonucleotides A, C, G, or T, the primer being at least 15 nucleotides in length, corresponding in sequence to the 3'-end of the bacteriophage-specific promoter, and extending across into at least the first two nucleotides of the cRNA, the mixture including all possibilities for the 3'-terminal two nucleotides;

(f) using the product of transcription in each of the sixteen subpools as a template for a polymerase chain reaction with a 3'-primer that corresponds in sequence to a sequence in the vector adjoining the site of insertion of the cDNA sample in the vector and a 5'-primer selected from the group consisting of: (i) the primer from which first-strand cDNA was made for that subpool; (ii) the primer from which the first-strand cDNA was made for that subpool extended at its 3'-terminus by an additional residue -N, where N can be any of A, C, G, or T; and (iii) the primer used for the synthesis of first-strand cDNA for that subpool extended at its 3'-terminus by two additional residues -N-N, wherein N can be any of A, C, G, or T, to produce polymerase chain reaction amplified fragments; and

(g) resolving the polymerase chain reaction amplified fragments by electrophoresis to display bands representing the 3'-ends of mRNAs present in the sample.

5

10

15

20

25

30

25

48

1900	1901	1902	1903	1904	1905	1906	1907	1908	1909	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921	1922	1923	1924	1925	1926	1927	1928	1929	1930	1931	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

11. The method of claim 1 wherein the first restriction endonuclease recognizing a four-nucleotide sequence is MspI.

5 12. The method of claim 1 wherein the first restriction endonuclease recognizing a four-nucleotide sequence is selected from the group consisting of TaqI and HinPII.

10 13. The method of claim 1 wherein the restriction endonuclease cleaving at a single site in each of the mixture of anchor primers is NotI.

15 14. The method of claim 1 wherein the step of generating linearized fragments of the cloned inserts comprises:

(i) dividing the plasmid containing the insert into two fractions, a first fraction cleaved with the restriction endonuclease XhoI and a second fraction cleaved with the restriction endonuclease SalI;

20 (ii) recombining the first and second fractions after cleavage;

25 (iii) dividing the recombined fractions into thirds and cleaving the first third with the restriction endonuclease HindIII, the second third with the restriction endonuclease BamHI, and the third third with the restriction endonuclease EcoRI; and

30 (iv) recombining the thirds after digestion in order to produce a population of linearized fragments of which about one-sixth of the population corresponds to the product of cleavage by each of the possible combinations of enzymes.

35 15. The method of claim 1 wherein the mRNA population has been enriched for polyadenylated mRNA species.

16. The method of claim 1 wherein the intensity of each band displayed after electrophoresis is about proportional to the abundance of the mRNA corresponding to the band in the original mixture.

17. The method of claim 16 further comprising a step of determining the relative abundance of each mRNA in the original mixture from the intensity of the band corresponding to that mRNA after electrophoresis.

18. The method of claim 1 wherein the step of resolving the polymerase chain reaction amplified fragments by electrophoresis comprises electrophoresis of the fragments on at least two gels.

19. The method of claim 1 wherein the suitable host cell is Escherichia coli.

20. The method of claim 1 further comprising the steps of:

(h) eluting at least one cDNA corresponding to a mRNA from an electropherogram in which bands representing the 3'-ends of mRNAs present in the sample are displayed;

(i) amplifying the eluted cDNA in a polymerase chain reaction;

(j) cloning the amplified cDNA into a plasmid;

(k) producing DNA corresponding to the cloned DNA from the plasmid; and

(l) sequencing the cloned cDNA.

21. A method for simultaneous sequence-specific identification of mRNAs in a mRNA population comprising the steps of:

(a) isolating a mRNA population;

(b) preparing double-stranded cDNAs from the mRNA population using a mixture of 12 anchor primers with the sequence A-A-C-T-G-G-A-A-G-A-A-T-T-C-G-C-G-G-C-C-G-C-A-G-G-A-A-T-T-T-T-T-T-T-T-T-T-T-T-T-T-T-T-V-N (SEQ ID NO: 2), wherein V is a deoxyribonucleotide selected from the group consisting of A, C, and G; and N is a deoxyribonucleotide selected from the group consisting of A, C, G, and T, the mixture including anchor primers containing all possibilities for V and N, to produce a cDNA sample;

(c) cleaving the cDNA sample with two restriction endonucleases, a first restriction endonuclease MspI and a second restriction endonuclease NotI;

(d) inserting the cDNA sample cleaved with the first and second restriction endonucleases into a vector, the cleaved cDNA being inserted in an orientation that is antisense with respect to a T3 promoter within the vector, the vector being the plasmid pBC SK⁺ cleaved with ClaI and NotI;

(e) transforming Escherichia coli with the vector into which the cleaved cDNA has been inserted to produce cloned inserts;

(f) generating linearized fragments of the cloned inserts by digestion with at least one restriction endonuclease that is different from the first and second restriction endonucleases;

(g) generating a cRNA preparation of antisense cRNA transcripts by incubation of the linearized fragments with a T3 RNA polymerase capable of initiating transcription from the T3-specific promoter;

(h) dividing the cRNA preparation into sixteen subpools and transcribing first-strand cDNA from each subpool, using a thermostable reverse transcriptase, and one of the sixteen primers A-G-G-T-C-G-A-C-G-G-T-A-T-C-G-G-N-N (SEQ ID NO: 3), wherein N is one of the four

deoxyribonucleotides A, C, G, or T, the mixture including all possibilities for the 3'-terminal two nucleotides;

(i) using the product of transcription in each of the sixteen subpools as a template for a polymerase chain reaction with the 3'-primer G-A-A-C-A-A-A-A-G-C-T-G-G-A-G-C-T-C-C-A-C-C-G-C (SEQ ID NO: 4), and a 5'-primer selected from the group consisting of: (1) the primer from which first-strand cDNA was made for that subpool; (2) the primer from which the first-strand cDNA was made for that subpool extended at its 3'-terminus by an additional residue -N, where N can be any of A, C, G, or T; and (3) the primer used for the synthesis of first-strand cDNA for that subpool extended at its 3'-terminus by two additional residues -N-N, wherein N can be any of A, C, G, or T, to produce polymerase chain reaction amplified fragments; and

(j) resolving the polymerase chain reaction amplified fragments by electrophoresis to display bands representing the 3'-ends of mRNAs present in the sample.

22. A method of simultaneous sequence-specific identification of mRNAs corresponding to members of an antisense cRNA pool representing the 3'-ends of a population of mRNAs, the antisense cRNAs that are members of the antisense cRNA pool being terminated at their 5'-end with a primer sequence corresponding to a bacteriophage-specific vector and at their 3'-end with a sequence corresponding in sequence to a sequence of the vector, the method comprising;

(a) dividing the members of the antisense cRNA pool into sixteen subpools and transcribing first-strand cDNA from each subpool, using a thermostable reverse transcriptase and one of sixteen primers whose 3'-terminus is -N-N, wherein N is one of the four deoxyribonucleotides A, C, G, or T, the primer being at least 15 nucleotides in length, corresponding in sequence

to the 3'-end of the bacteriophage-specific promoter, and extending across into at least the first two nucleotides of the cRNA, the mixture including all possibilities for the 3'-terminal two nucleotides;

5 (b) using the product of transcription in each of the sixteen subpools as a template for a polymerase chain reaction with a 3'-primer that corresponds in sequence to a sequence vector adjoining the site of insertion of the cDNA sample in the vector and a 5'-
10 primer selected from the group consisting of: (i) the primer from which first-strand cDNA was made for that subpool; (ii) the primer from which the first-strand cDNA was made for that subpool extended at its 3'-terminus by an additional residue -N, where N can be any of A, C, G,
15 or T; and (iii) the primer used for the synthesis of first-strand cDNA for that subpool extended at its 3'-terminus by two additional residues -N-N, wherein N can be any of A, C, G, or T, to produce polymerase chain reaction amplified fragments; and

20 (c) resolving the polymerase chain reaction amplified fragments by electrophoresis to display bands representing the 3'-ends of mRNAs present in the sample.

25 23. A method for detecting a change in the pattern of mRNA expression in a tissue associated with a physiological or pathological change comprising the steps of:

30 (a) obtaining a first sample of a tissue that is not subject to the physiological or pathological change;

35 (b) determining the pattern of mRNA expression in the first sample of the tissue by performing steps (a)-(c) of claim 22 to generate a first display of bands representing the 3'-ends of mRNAs present in the first sample;

(c) obtaining a second sample of the tissue that has been subject to the physiological or pathological change;

5 (d) determining the pattern of mRNA expression in the second sample of the tissue by performing steps (a)-(c) of claim 22 to generate a second display of bands representing the 3'-ends of mRNAs present in the second sample; and

10 (e) comparing the first and second displays to determine the effect of the physiological or pathological change on the pattern of mRNA expression in the tissue.

15 24. The method of claim 23 wherein the tissue is derived from the central nervous system.

20 25. The method of claim 24 wherein the physiological or pathological change is selected from the group consisting of Alzheimer's disease, parkinsonism, ischemia, alcohol addiction, drug addiction, schizophrenia, amyotrophic lateral sclerosis, multiple sclerosis, depression, and bipolar manic-depressive disorder.

5 26. The method of claim 24 wherein the physiological or pathological change is associated with learning or memory, emotion, glutamate neurotoxicity, feeding behavior, olfaction, vision, movement disorders, viral infection, electroshock therapy, or the administration of a drug or toxin.

30 27. The method of claim 24 wherein the physiological or pathological change is selected from the group consisting of circadian variation, aging, and long-term potentiation.

28. The method of claim 24 wherein the tissue is derived from a structure within the central nervous system selected from the group consisting of retina, cerebral cortex, olfactory bulb, thalamus, hypothalamus, anterior pituitary, posterior pituitary, hippocampus, nucleus accumbens, amygdala, striatum, cerebellum, brain stem, suprachiasmatic nucleus, and spinal cord.

29. The method of claim 23 wherein the tissue is from an organ or organ system selected from the group consisting of the cardiovascular system, the pulmonary system, the digestive system, the peripheral nervous system, the liver, the kidney, skeletal muscle, and the reproductive system.

30. A method of screening for a side effect of a drug comprising the steps of:

(a) obtaining a first sample of tissue from an organism treated with a compound of known physiological function;

(b) determining the pattern of mRNA expression in the first sample of the tissue by performing steps (a)-(c) of claim 22 to generate a first display of bands representing the 3'-ends of mRNAs present in the first sample;

(c) obtaining a second sample of tissue from an organism treated with a drug to be screened for a side effect;

(d) determining the pattern of mRNA expression in the second sample of the tissue by performing steps (a)-(c) of claim 22 to generate a second display of bands representing the 3'-ends of mRNAs present in the second sample; and

(e) comparing the first and second displays in order to detect the presence of mRNA species whose expression is not affected by the known compound but is

affected by the drug to be screened, thereby indicating a difference in action of the drug to be screened and the known compound and thus a side effect.

31. The method of claim 30 wherein the drug to be tested is selected from the group consisting of antidepressants, neuroleptics, tranquilizers, anticonvulsants, monoamine oxidase inhibitors, and stimulants.

32. The method of claim 30 wherein the drug to be tested is selected from the group consisting of anti-parkinsonism agents, skeletal muscle relaxants, analgesics, local anesthetics, cholinergics, antiviral agents, antispasmodics, steroids, and non-steroidal anti-inflammatory drugs.

33. A panel of primers comprising 16 primers of the sequence A-G-G-T-C-G-A-C-G-G-T-A-T-C-G-G-N-N (SEQ ID NO: 3), wherein N is one of the four deoxyribonucleotides A, C, G, or T.

34. A panel of primers comprising 64 primers of the sequences A-G-G-T-C-G-A-C-G-G-T-A-T-C-G-G-N-N-N (SEQ ID NO: 5), wherein N is one of the four deoxyribonucleotides A, C, G, or T.

35. A panel of primers comprising 256 primers of the sequences A-G-G-T-C-G-A-C-G-G-T-A-T-C-G-G-N-N-N-N (SEQ ID NO: 6), wherein N is one of the four deoxyribonucleotides A, C, G, or T.

36. A panel of primers comprising 12 primers of the sequences A-A-C-T-G-G-A-A-G-A-A-T-T-C-G-C-G-G-C-C-G-C-A-G-G-A-A-T-T-T-T-T-T-T-T-T-T-T-T-T-T-T-T-T-V-N (SEQ ID NO: 2) wherein V is a deoxyribonucleotide

selected from the group consisting of A, C, and G; and N is a deoxyribonucleotide selected from the group consisting of A, C, G, and T.

5 37. A degenerate mixture of primers comprising
a mixture of 12 primers of the sequences A-A-C-T-G-G-A-A-
G-A-A-T-T-C-G-C-G-G-C-C-G-C-A-G-G-A-A-T-T-T-T-T-T-T-T-
T-T-T-T-T-T-T-T-T-V-N (SEQ ID NO: 2), wherein V is a
10 deoxyribonucleotide selected from the group consisting of
A, C, and G; and N is a deoxyribonucleotide selected from
the group consisting of A, C, G, and T, each of the 12
primers being present in about an equimolar quantity.